

### **AMENDMENTS TO THE CLAIMS**

Please amend the claims as follows:

#### **LISTING OF CLAIMS:**

1. (Currently Amended): A process for the production of L-ascorbic acid comprising:

(a) contacting an enzyme with a substrate which is selected from the group consisting of L-gulose, L-galactose, L-idose, and L-talose;

(b) converting the substrate directly into L-ascorbic acid by catalytic activity of the enzyme under suitable culture conditions; and

(c) isolating L-ascorbic acid from the reaction mixture,

wherein said enzyme has (1) the amino acid sequence of SEQ ID NO: 2 or (2) ~~an amino acid sequence with 90% sequence identity to SEQ ID NO: 2 and with the activity to produce L-ascorbic acid or~~ (3) an amino acid sequence encoded by the DNA sequence of SEQ ID NO: 1 or (4) ~~(3)~~ (3) an amino acid sequence encoded by a DNA sequence that hybridizes to the complement of DNA sequence of SEQ ID NO: 1 under highly stringent hybridization conditions of 6X SSC, 0.5% SDS, 100 µg/ml denatured salmon sperm DNA, 50% formamide, and incubating overnight at 42°C with gentle rocking and highly stringent wash conditions of washing in 2X SSC, 0.5% SDS at room temperature for 15 minutes, followed by another wash in 0.1X SSC, 0.5% SDS at room temperature for 15 minutes ~~to the DNA sequence of SEQ ID: 1~~ and having the activity to produce L-ascorbic acid.

2. (Currently Amended): A process for the production of L-ascorbic acid with an enzyme having (1) the amino acid sequence of SEQ ID NO: 2 or (2) ~~an amino acid sequence with 90% sequence identity to SEQ ID NO: 2 and with the activity to produce L-ascorbic acid or~~ (3) an amino acid sequence encoded by the DNA sequence of SEQ ID NO: 1 or (4) ~~(3)~~ (3) an amino acid sequence encoded by a DNA sequence that hybridizes to the complement of DNA sequence of SEQ ID NO: 1 under highly stringent hybridization conditions of 6X SSC, 0.5% SDS, 100 µg/ml denatured salmon sperm DNA, 50% formamide, and incubating overnight at 42°C with gentle rocking and highly stringent wash conditions of washing in 2X SSC, 0.5% SDS at room temperature for 15 minutes, followed by another wash in 0.1X SSC, 0.5% SDS at room temperature for 15 minutes ~~to the DNA sequence of SEQ ID: 1~~ and having the activity to produce L-ascorbic acid, whereby L-ascorbic acid is produced from a substrate which is selected from the group consisting of L-gulono-1,4-lactone, L-gulonic acid, L-galactono-1,4-lactone, L-galactonic acid, L-idono-1,4-lactone, L-idonic acid, L-talono-1,4-lactone, and L-talonic acid,

said process comprising the steps of:

- (a) contacting the enzyme with the substrate,
- (b) converting the substrate directly into L-ascorbic acid by catalytic activity of the enzyme under suitable culture conditions; and
- (c) isolating L-ascorbic acid from the reaction mixture.

3. (Withdrawn): A process for the production of L-gulono-1,4-lactone or L-gulonic acid with an enzyme having the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence that is 90% identical thereto, with the activity to produce L-gulono-

1,4-lactone or L-gulonic acid, whereby L-gulono-1,4-lactone or L-gulonic acid is produced from L-gulose.

4. (Withdrawn): A process for the production of L-galactono-1,4-lactone or L-galactonic acid with an enzyme having the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence that is 90% identical thereto, with the activity to produce L-galactono-1,4-lactone or L-galactonic acid, whereby L-galactono-1,4-lactone or L-galactonic acid is produced from L-galactose.

5. (Cancelled).

6. (Previously presented): A process according to claim 1, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

7. (Original): A process according to claim 6, wherein the process is conducted at a pH of about 2 to about 8 and at a temperature of about 18°C to about 42°C.

8. (Previously presented): A process for producing L-ascorbic acid comprising (a) contacting a substrate which is selected from the group consisting of L-gulose, L-galactose, L-idose, L-talose, L-gulono-1,4-lactone, L-gulonic acid, L-galactono-1,4-lactone, and L-galactonic acid with an enzyme derivable from *G. oxydans* DSM 4025, (b) converting the substrate directly into L-ascorbic acid by catalytic activity of the enzyme under suitable culture conditions and (c) isolating L-ascorbic acid from the reaction mixture; wherein the enzyme has the following physico-chemical properties:

(a) molecular weight of about 60,000 Da on SDS-PAGE;

(b) substrate specificity for primary and secondary alcohols and aldehydes;

(c) pH-stability at pH of about 6 to about 9;

(d) pH-optimum at pH of about 8.0; and

(e) inhibited by  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$ , and  $\text{Fe}^{3+}$ .

9. (Withdrawn): A process for producing L-gulono-1,4-lactone or L-gulonic acid comprising contacting L-gulose with Enzyme B of *G. oxydans* DSM 4025 and isolating L-gulono-1,4-lactone or L-gulonic acid from the reaction mixture, wherein Enzyme B has the following physico-chemical properties:

(a) molecular weight of about 60,000 Da on SDS-PAGE;

(b) substrate specificity for primary and secondary alcohols and aldehydes;

(c) pH-stability at pH of about 6 to about 9;

(d) pH-optimum at pH of about 8.0; and

(e) inhibited by  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$ , and  $\text{Fe}^{3+}$ .

10. (Withdrawn): A process for producing L-galactono-1,4-lactone or galactonic acid comprising contacting L-galactose with Enzyme B of *G. oxydans* DSM 4025 and isolating L-galactono-1,4-lactone or galactonic acid from the reaction mixture, wherein Enzyme B has the following physico-chemical properties:

(a) molecular weight of about 60,000 Da on SDS-PAGE;

(b) substrate specificity for primary and secondary alcohols and aldehydes;

(c) pH-stability at pH of about 6 to about 9;

(d) pH-optimum at pH of about 8.0; and

(e) inhibited by  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$ , and  $\text{Fe}^{3+}$ .

11. (Withdrawn): A process according to claim 3 comprising (a) contacting the enzyme with the substrate and (b) isolating the L-gulono-1,4-lactone or L-gulonic acid from the reaction mixture.

12. (Withdrawn): A process according to claim 4 comprising (a) contacting the enzyme with the substrate and (b) isolating the L-galactono-1,4-lactone or L-galactonic acid from the reaction mixture.

13. (Previously presented): A process according to claim 2, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

14. (Withdrawn): A process according to claim 3, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

15. (Withdrawn): A process according to claim 4, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

16. (Previously presented): A process according to claim 8, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

17. (Withdrawn): A process according to claim 11, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

Application No.: 10/528,673

Amendment Dated: August 1, 2008

Reply to Office Action Dated: February 1, 2008

18. (Withdrawn): A process according to claim 12, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.